arXiv:0908.0097v2 [math.DG] 8 Aug 2009

Multi-Time KCC-Invariants

Mircea Neagu

1 August 2009; Revised 7 August 2009 (minor corrections)

Abstract

In this paper we construct some multi-time geometrical extensions of the KCC-invariants, which characterize a given second-order system of PDEs on the 1-jet space $J^1(T,M)$. A theorem of characterization of these multi-time geometrical KCC-invariants is given.

Mathematics Subject Classification (2000): 58A20, 58B20, 35M99.

Key words and phrases: multi-time 1-jet spaces, temporal and spatial semisprays, nonlinear connections, PDE systems of second order, multi-time KCC-invariants.

1 Geometrical objects on multi-time 1-jet spaces

We remind first few differential geometrical properties of the multi-time 1-jet spaces. The multi-time 1-jet bundle

$$\xi_1 = (J^1(T,M), \pi_1, T \times M)$$

is a vector bundle over the product manifold $T \times M$, having the fibre type $\mathbb{R}^{mn}$, where m (resp. n) is the dimension of the *temporal* (resp. *spatial*) manifold T (resp. M). If the temporal manifold T has the local coordinates $(t^\alpha)_{\alpha=\overline{1,m}}$ and the spatial manifold M has the local coordinates $(x^i)_{i=\overline{1,n}}$, then we denote the local coordinates of the multi-time 1-jet space $J^1(T,M)$ by $(t^\alpha, x^i, x^i_\alpha)$. These transform by the rules [8]

$$\begin{cases} \widetilde{t}^\alpha = \widetilde{t}^\alpha(t^\beta) \\ \widetilde{x}^i = \widetilde{x}^i(x^j) \\ \widetilde{x}^i_\alpha = \dfrac{\partial \widetilde{x}^i}{\partial x^j} \dfrac{\partial t^\beta}{\partial \widetilde{t}^\alpha} x^j_\beta, \end{cases} \tag{1.1}$$

where $\det(\partial \widetilde{t}^\alpha / \partial t^\beta) \neq 0$ and $\det(\partial \widetilde{x}^i / \partial x^j) \neq 0$.

Remark 1.1 *In this work the greek indices* $\alpha, \beta, \gamma, \delta, \varepsilon, \mu, \nu, \rho$... *run over the set* $\{1, 2, ..., m\}$ *and the latin indices* i, j, k, l, p, q, r, s... *run over the set* $\{1, 2, ..., n\}$. *The Einstein convention of summation is also adopted all over this paper.*

In the geometrical study of the multi-time 1-jet vector bundle, a central rôle is played by the *distinguished tensors* ($d-$tensors).

Definition 1.2 *A geometrical object* $D = \left(D^{\alpha i(j)(\nu)\ldots}_{\gamma k(\beta)(l)\ldots}\right)$ *on the 1-jet vector bundle* $J^1(T,M)$*, whose local components transform by the rules*

$$D^{\alpha i(j)(\nu)\ldots}_{\gamma k(\beta)(l)\ldots} = \widetilde{D}^{\delta p(q)(\eta)\ldots}_{\varepsilon r(\mu)(s)\ldots} \frac{\partial t^\alpha}{\partial \widetilde{t}^\delta} \frac{\partial x^i}{\partial \widetilde{x}^p} \left(\frac{\partial x^j}{\partial \widetilde{x}^q} \frac{\partial \widetilde{t}^\mu}{\partial t^\beta}\right) \frac{\partial \widetilde{t}^\varepsilon}{\partial t^\gamma} \frac{\partial \widetilde{x}^r}{\partial x^k} \left(\frac{\partial \widetilde{x}^s}{\partial x^l} \frac{\partial t^\nu}{\partial \widetilde{t}^\eta}\right) \ldots, \tag{1.2}$$

is called a $d-$***tensor field****.*

Remark 1.3 *The use of parentheses for certain indices of the local components* $D^{\alpha i(j)(\nu)\ldots}_{\gamma k(\beta)(l)\ldots}$ *of the distinguished tensor field* D *on the 1-jet space is motivated by the fact that the pair of indices "* $\substack{(j)\\(\beta)}$ *" or "* $\substack{(\nu)\\(l)}$ *" behaves like a single index.*

Example 1.4 *The geometrical object*

$$\mathbf{C} = \mathbf{C}^{(i)}_{(\alpha)} \frac{\partial}{\partial x^i_\alpha},$$

where $\mathbf{C}^{(i)}_{(\alpha)} = x^i_\alpha$*, represents a* $d-$*tensor field on the 1-jet space; this is called the* ***canonical Liouville*** $d-$***tensor field*** *of the 1-jet vector bundle* $J^1(T,M)$ *and it is a global geometrical object.*

Example 1.5 *Let* $h = (h_{\alpha\beta}(t))$ *be a Riemannian metric on the temporal manifold* T*. The geometrical object*

$$\mathbf{J}_h = J^{(i)}_{(\alpha)\beta j} \frac{\partial}{\partial x^i_\alpha} \otimes dt^\beta \otimes dx^j,$$

where $J^{(i)}_{(\alpha)\beta j} = h_{\alpha\beta}\delta^i_j$ *is a* $d-$*tensor field on* $J^1(T,M)$*, which is called the* ***h-normalization*** $d-$***tensor field*** *of the 1-jet space* $J^1(T,M)$*. Obviously, it is also a global geometrical object.*

In the Riemann-Lagrange differential geometry of the 1-jet spaces developed in [8], [9] important rôles are also played by geometrical objects as the *temporal* or *spatial semisprays*, together with the *multi-time jet nonlinear connections.*

Definition 1.6 *A set of local functions* $H = \left(H^{(i)}_{(\alpha)\beta}\right)$ *on* $J^1(T,M)$*, which transform by the rules*

$$2\widetilde{H}^{(i)}_{(\alpha)\beta} = 2H^{(k)}_{(\gamma)\nu} \frac{\partial \widetilde{x}^i}{\partial x^k} \frac{\partial t^\gamma}{\partial \widetilde{t}^\alpha} \frac{\partial t^\nu}{\partial \widetilde{t}^\beta} - \frac{\partial t^\mu}{\partial \widetilde{t}^\beta} \frac{\partial \widetilde{x}^i_\alpha}{\partial t^\mu}, \tag{1.3}$$

is called a ***temporal semispray*** *on* $J^1(T,M)$*.*

Example 1.7 *Let us consider a Riemannian metric $h = (h_{\alpha\beta}(t))$ on the temporal manifold T and let*

$$H^{\alpha}_{\beta\gamma} = \frac{h^{\alpha\mu}}{2}\left(\frac{\partial h_{\beta\mu}}{\partial t^{\gamma}} + \frac{\partial h_{\gamma\mu}}{\partial t^{\beta}} - \frac{\partial h_{\beta\gamma}}{\partial t^{\mu}}\right)$$

be its Christoffel symbols. Taking into account that we have the transformation rules

$$\widetilde{H}^{\delta}_{\nu\rho} = H^{\alpha}_{\beta\gamma}\frac{\partial \widetilde{t}^{\delta}}{\partial t^{\alpha}}\frac{\partial t^{\beta}}{\partial \widetilde{t}^{\nu}}\frac{\partial t^{\gamma}}{\partial \widetilde{t}^{\rho}} + \frac{\partial \widetilde{t}^{\delta}}{\partial t^{\varepsilon}}\frac{\partial^2 t^{\varepsilon}}{\partial \widetilde{t}^{\nu}\partial \widetilde{t}^{\rho}}, \tag{1.4}$$

we deduce that the local components

$$\mathring{H}^{(i)}_{(\alpha)\beta} = -\frac{1}{2}H^{\mu}_{\alpha\beta}x^{i}_{\mu}$$

define a temporal semispray $\mathring{H} = \left(\mathring{H}^{(i)}_{(\alpha)\beta}\right)$ on $J^1(T, M)$. This is called the ***canonical temporal semispray associated to the temporal metric*** *$h_{\alpha\beta}(t)$.*

Definition 1.8 *A set of local functions $G = \left(G^{(i)}_{(\alpha)\beta}\right)$, which transform by the rules*

$$2\widetilde{G}^{(i)}_{(\alpha)\beta} = 2G^{(k)}_{(\gamma)\nu}\frac{\partial \widetilde{x}^{i}}{\partial x^{k}}\frac{\partial t^{\gamma}}{\partial \widetilde{t}^{\alpha}}\frac{\partial t^{\nu}}{\partial \widetilde{t}^{\beta}} - \frac{\partial x^{r}}{\partial \widetilde{x}^{s}}\frac{\partial \widetilde{x}^{i}_{\alpha}}{\partial x^{r}}\widetilde{x}^{s}_{\beta}, \tag{1.5}$$

is called a ***spatial semispray*** *on $J^1(T, M)$.*

Example 1.9 *Let $\varphi = (\varphi_{ij}(x))$ be a Riemannian metric on the spatial manifold M and let us consider*

$$\gamma^{i}_{jk} = \frac{\varphi^{im}}{2}\left(\frac{\partial \varphi_{jm}}{\partial x^{k}} + \frac{\partial \varphi_{km}}{\partial x^{j}} - \frac{\partial \varphi_{jk}}{\partial x^{m}}\right)$$

its Christoffel symbols. Taking into account that we have the transformation rules

$$\widetilde{\gamma}^{p}_{qr} = \gamma^{i}_{jk}\frac{\partial \widetilde{x}^{p}}{\partial x^{i}}\frac{\partial x^{j}}{\partial \widetilde{x}^{q}}\frac{\partial x^{k}}{\partial \widetilde{x}^{r}} + \frac{\partial \widetilde{x}^{p}}{\partial x^{l}}\frac{\partial^2 x^{l}}{\partial \widetilde{x}^{q}\partial \widetilde{x}^{r}}, \tag{1.6}$$

we deduce that the local components

$$\mathring{G}^{(i)}_{(\alpha)\beta} = \frac{1}{2}\gamma^{i}_{pq}x^{p}_{\alpha}x^{q}_{\beta}$$

define a spatial semispray $\mathring{G} = \left(\mathring{G}^{(i)}_{(\alpha)\beta}\right)$ on $J^1(T, M)$. This is called the ***canonical spatial semispray associated to the spatial metric*** *$\varphi_{ij}(x)$.*

Definition 1.10 *A set of local functions $\Gamma = \left(M^{(i)}_{(\alpha)\beta}, N^{(i)}_{(\alpha)j}\right)$ on $J^1(T, M)$, which transform by the rules*

$$\widetilde{M}^{(i)}_{(\alpha)\beta} = M^{(k)}_{(\gamma)\nu}\frac{\partial \widetilde{x}^{i}}{\partial x^{k}}\frac{\partial t^{\gamma}}{\partial \widetilde{t}^{\alpha}}\frac{\partial t^{\nu}}{\partial \widetilde{t}^{\beta}} - \frac{\partial t^{\mu}}{\partial \widetilde{t}^{\beta}}\frac{\partial \widetilde{x}^{i}_{\alpha}}{\partial t^{\mu}} \tag{1.7}$$

and

$$\widetilde{N}^{(i)}_{(\alpha)j} = N^{(k)}_{(\gamma)l}\frac{\partial \widetilde{x}^i}{\partial x^k}\frac{\partial t^\gamma}{\partial \widetilde{t}^\alpha}\frac{\partial x^l}{\partial \widetilde{x}^j} - \frac{\partial x^r}{\partial \widetilde{x}^j}\frac{\partial \widetilde{x}^i_\alpha}{\partial x^r}, \tag{1.8}$$

is called a ***nonlinear connection*** *on the 1-jet space* $J^1(T,M)$.

Example 1.11 *Let us consider that* $(T, h_{\alpha\beta}(t))$ *and* $(M, \varphi_{ij}(x))$ *are Riemannian manifolds having the Christoffel symbols* $H^\alpha_{\beta\gamma}(t)$ *and* $\gamma^i_{jk}(x)$*. Then, using the transformation rules (1.1), (1.4) and (1.6), we deduce that the set of local functions*

$$\mathring{\Gamma} = \left(\mathring{M}^{(i)}_{(\alpha)\beta}, \mathring{N}^{(i)}_{(\alpha)j}\right),$$

where

$$\mathring{M}^{(i)}_{(\alpha)\beta} = -H^\mu_{\alpha\beta}x^i_\mu \quad \text{and} \quad \mathring{N}^{(i)}_{(\alpha)j} = \gamma^i_{jr}x^r_\alpha,$$

represents a nonlinear connection on the 1-jet space $J^1(T,M)$*. This multi-time jet nonlinear connection is called the* ***canonical nonlinear connection attached to the pair of Riemannian metrics*** $(h_{\alpha\beta}(t), \varphi_{ij}(x))$.

In the sequel, let us study the geometrical relations between *temporal* or *spatial semisprays* and *multi-time nonlinear connections* on the 1-jet vector bundle $J^1(T,M)$. In this direction, using the local transformation laws (1.3), (1.7) and (1.1), respectively the transformation laws (1.5), (1.8) and (1.1), by direct local computations, we find the following geometrical results:

Theorem 1.12 *The temporal semisprays* $H = \left(H^{(i)}_{(\alpha)\beta}\right)$ *and the sets of temporal components of nonlinear connections* $\Gamma_{temporal} = \left(M^{(i)}_{(\alpha)\beta}\right)$ *are in one-to-one correspondence on the 1-jet space* $J^1(T,M)$*, via:*

$$M^{(i)}_{(\alpha)\beta} = 2H^{(i)}_{(\alpha)\beta}, \qquad H^{(i)}_{(\alpha)\beta} = \frac{1}{2}M^{(i)}_{(\alpha)\beta};$$

Theorem 1.13 ***(i)*** *If* $G^{(i)}_{(\alpha)\beta}$ *are the components of a spatial semispray on* $J^1(T,M)$*, where* $(T, h_{\alpha\beta}(t))$ *is a Riemannian manifold, then the components*

$$N^{(i)}_{(\alpha)j} = \frac{\partial G^i}{\partial x^j_\mu}h_{\alpha\mu},$$

where $G^i = h^{\delta\varepsilon}G^{(i)}_{(\delta)\varepsilon}$*, represent a spatial nonlinear connection on* $J^1(T,M)$*.*

(ii) *Conversely, the spatial nonlinear connection* $\Gamma_{spatial} = \left(N^{(i)}_{(\alpha)j}\right)$ *produces the spatial semispray components*

$$G^{(i)}_{(\alpha)\beta} = \frac{1}{2}N^{(i)}_{(\alpha)r}x^r_\beta.$$

2 Multi-time geometrical KCC-theory

In this Section we construct some multi-time generalizations on the 1-jet space $J^1(T, M)$ for the basic objects of the KCC-theory ([1], [2], [3], [10]). In this respect, let us consider on $J^1(T, M)$ a second-order system of partial differential equations of local form

$$\frac{\partial^2 x^i}{\partial t^\alpha \partial t^\beta} + F^{(i)}_{(\alpha)\beta}(t^\gamma, x^k, x^k_\gamma) = 0, \quad \alpha, \beta = \overline{1, m}, \quad i = \overline{1, n}, \tag{2.1}$$

where $x^k_\gamma = \partial x^k / \partial t^\gamma$, $F^{(i)}_{(\alpha)\beta} = F^{(i)}_{(\beta)\alpha}$ and the local components $F^{(i)}_{(\alpha)\beta}(t^\gamma, x^k, x^k_\gamma)$ transform under a change of coordinates (1.1) by the rules

$$\widetilde{F}^{(i)}_{(\alpha)\beta} = 2F^{(k)}_{(\gamma)\nu} \frac{\partial \widetilde{x}^i}{\partial x^k} \frac{\partial t^\gamma}{\partial \widetilde{t}^\alpha} \frac{\partial t^\nu}{\partial \widetilde{t}^\beta} - \frac{\partial t^\mu}{\partial \widetilde{t}^\beta} \frac{\partial \widetilde{x}^i_\alpha}{\partial t^\mu} - \frac{\partial x^r}{\partial \widetilde{x}^s} \frac{\partial \widetilde{x}^i_\alpha}{\partial x^r} \widetilde{x}^s_\beta, . \tag{2.2}$$

Remark 2.1 *The second-order system of partial differential equations (2.1) is invariant under a change of coordinates (1.1).*

Example 2.2 *Let us consider that $(T, h_{\alpha\beta}(t))$ and $(M, \varphi_{ij}(x))$ are Riemannian manifolds having the Christoffel symbols $H^\alpha_{\beta\gamma}(t)$ and $\gamma^i_{jk}(x)$. Then, the local components*

$$\overset{\circ}{F}{}^{(i)}_{(\alpha)\beta} = -H^\mu_{\alpha\beta} x^i_\mu + \gamma^i_{pq} x^p_\alpha x^q_\beta$$

transform under a change of coordinates (1.1) by the rules (2.2). In this particular case, the PDE system (2.1) becomes

$$\frac{\partial^2 x^i}{\partial t^\alpha \partial t^\beta} - H^\mu_{\alpha\beta} x^i_\mu + \gamma^i_{pq} x^p_\alpha x^q_\beta = 0, \quad \alpha, \beta = \overline{1, m}, \quad i = \overline{1, n}, \tag{2.3}$$

that is the PDE system of the affine maps between the Riemannian manifolds $(T, h_{\alpha\beta}(t))$ and $(M, \varphi_{ij}(x))$. We recall that these affine maps carry the geodesics of the temporal Riemannian manifold $(T, h_{\alpha\beta}(t))$ into the geodesics of the spatial Riemannian manifold $(M, \varphi_{ij}(x))$. Moreover, the $h-$trace of the equations (2.3) produces the equations of the harmonic maps between the Riemannian manifolds $(T, h_{\alpha\beta}(t))$ and $(M, \varphi_{ij}(x))$. For more details, please see [6].

Using a temporal Riemannian metric $h_{\alpha\beta}(t)$ on T and taking into account the transformation rules (1.3), (1.5) and (2.2), we can rewrite the PDE system (2.1) in the following form:

$$\frac{\partial^2 x^i}{\partial t^\alpha \partial t^\beta} - H^\mu_{\alpha\beta} x^i_\mu + 2G^{(i)}_{(\alpha)\beta}(t^\gamma, x^k, x^k_\gamma) = 0, \quad \alpha, \beta = \overline{1, m}, \quad i = \overline{1, n},$$

where

$$G^{(i)}_{(\alpha)\beta} = \frac{1}{2} F^{(i)}_{(\alpha)\beta} + \frac{1}{2} H^\mu_{\alpha\beta} x^i_\mu$$

are the components of a spatial semispray on $J^1(T, M)$. The coefficients of the spatial semispray $G^{(i)}_{(\alpha)\beta}$ produce the spatial components $N^{(i)}_{(\alpha)j}$ of a nonlinear connection Γ on the 1-jet space $J^1(T, M)$, by putting

$$N^{(i)}_{(\alpha)j} = \frac{h^{\mu\nu}\partial G^{(i)}_{(\mu)\nu}}{\partial x^j_\gamma} h_{\gamma\alpha} = \frac{h^{\mu\nu}}{2}\frac{\partial F^{(i)}_{(\mu)\nu}}{\partial x^j_\gamma} h_{\gamma\alpha} + \frac{h^{\mu\nu}}{2} H^\gamma_{\mu\nu} h_{\gamma\alpha}\delta^i_j.$$

In order to find the basic jet multi-time geometrical invariants of the PDE system (2.1) (please see Kosambi [7], Cartan [4] and Chern [5]) under the coordinate transformations (1.1), we define the *$h-$KCC-covariant derivative of a $d-$tensor of type* $T^{(i)}_{(\alpha)}(t^\gamma, x^k, x^k_\gamma)$ on the 1-jet space $J^1(T, M)$, via

$$\begin{aligned}\frac{\overset{h}{\nabla} T^{(i)}_{(\alpha)}}{\partial t^\beta} &= \frac{\partial T^{(i)}_{(\alpha)}}{\partial t^\beta} + N^{(i)}_{(\alpha)r} T^{(r)}_{(\beta)} - H^\mu_{\alpha\beta} T^{(i)}_{(\mu)} = \\ &= \frac{\partial T^{(i)}_{(\alpha)}}{\partial t^\beta} + \frac{h^{\mu\nu}}{2}\frac{\partial F^{(i)}_{(\mu)\nu}}{\partial x^r_\gamma} h_{\gamma\alpha} T^{(r)}_{(\beta)} + \frac{h^{\mu\nu}}{2} H^\gamma_{\mu\nu} h_{\gamma\alpha} T^{(i)}_{(\beta)} - H^\mu_{\alpha\beta} T^{(i)}_{(\mu)}.\end{aligned}$$

Remark 2.3 *The $h-$KCC-covariant derivative components* $\frac{\overset{h}{\nabla} T^{(i)}_{(\alpha)}}{\partial t^\beta}$ *transform under a change of coordinates (1.1) as a $d-$tensor of type $\mathcal{T}^{(i)}_{(\alpha)\beta}$.*

In such a geometrical context, if we use the notation $x^i_\alpha = \partial x^i / \partial t^\alpha$, then the PDE system (2.1) can be rewritten in the following distinguished tensorial form:

$$\begin{aligned}\frac{\overset{h}{\nabla} x^i_\alpha}{\partial t^\beta} &= -F^{(i)}_{(\alpha)\beta}(t^\gamma, x^k, x^k_\gamma) + N^{(i)}_{(\alpha)r} x^r_\beta - H^\mu_{\alpha\beta} x^i_\mu = \\ &= -F^{(i)}_{(\alpha)\beta} + \frac{h^{\mu\nu}}{2}\frac{\partial F^{(i)}_{(\mu)\nu}}{\partial x^r_\gamma} h_{\gamma\alpha} x^r_\beta + \frac{h^{\mu\nu}}{2} H^\gamma_{\mu\nu} h_{\gamma\alpha} x^i_\beta - H^\mu_{\alpha\beta} x^i_\mu.\end{aligned}$$

Definition 2.4 *The distinguished tensor*

$$\overset{h}{\varepsilon}{}^{(i)}_{(\alpha)\beta} = -F^{(i)}_{(\alpha)\beta} + \frac{h^{\mu\nu}}{2}\frac{\partial F^{(i)}_{(\mu)\nu}}{\partial x^r_\gamma} h_{\gamma\alpha} x^r_\beta + \frac{h^{\mu\nu}}{2} H^\gamma_{\mu\nu} h_{\gamma\alpha} x^i_\beta - H^\mu_{\alpha\beta} x^i_\mu$$

*is called the **first multi-time** $h-$**KCC-invariant** on the 1-jet space $J^1(T, M)$ of the PDEs (2.1), which can be interpreted as an **external force** [1], [3].*

Example 2.5 *For the second-order PDE system (2.3), which gives the affine maps between the Riemannian manifolds $(T, h_{\alpha\beta}(t))$ and $(M, \varphi_{ij}(x))$, the first multi-time $h-$KCC-invariant is zero.*

Example 2.6 *It can be easily seen that for the particular first order PDE system*

$$\frac{\partial x^i}{\partial t^\alpha} = X^{(i)}_{(\alpha)}(t^\gamma, x^k) \Rightarrow \frac{\partial^2 x^i}{\partial t^\alpha \partial t^\beta} = \frac{\partial X^{(i)}_{(\alpha)}}{\partial t^\beta} + \frac{\partial X^{(i)}_{(\alpha)}}{\partial x^r} x^r_\beta, \tag{2.4}$$

where $X^{(i)}_{(\alpha)}(t,x)$ is a given $d-$tensor on $J^1(T,M)$, the first multi-time $h-$KCC-invariant has the form

$$\overset{h}{\varepsilon}{}^{(i)}_{(\alpha)\beta} = \frac{\partial X^{(i)}_{(\alpha)}}{\partial t^\beta} + \frac{1}{2}\frac{\partial X^{(i)}_{(\alpha)}}{\partial x^r} x^r_\beta + \frac{h^{\mu\nu}}{2} H^\gamma_{\mu\nu} h_{\gamma\alpha} x^i_\beta - H^\mu_{\alpha\beta} x^i_\mu.$$

In the sequel, let us vary the solutions $x^i(t^\gamma)$ of the PDE system (2.1) by the nearby smooth maps $(\overline{x}^i(t^\gamma, s))_{s\in(-\varepsilon,\varepsilon)}$, where $\overline{x}^i(t^\gamma, 0) = x^i(t^\gamma)$. Then, if we consider the *variation $d-$tensor field*

$$\xi^i(t^\gamma) = \left.\frac{\partial \overline{x}^i}{\partial s}\right|_{s=0},$$

we get the *variational equations*

$$\frac{\partial^2 \xi^i}{\partial t^\alpha \partial t^\beta} + \frac{\partial F^{(i)}_{(\alpha)\beta}}{\partial x^k}\xi^k + \frac{\partial F^{(i)}_{(\alpha)\beta}}{\partial x^r_\mu}\frac{\partial \xi^r}{\partial t^\mu} = 0, \quad \alpha, \beta = \overline{1,m}, \quad i = \overline{1,n}. \tag{2.5}$$

It is obvious that the equations (2.5) imply the *$h-$trace variational equations*

$$h^{\alpha\beta}\frac{\partial^2 \xi^i}{\partial t^\alpha \partial t^\beta} + \frac{\partial F^i}{\partial x^k}\xi^k + \frac{\partial F^i}{\partial x^r_\mu}\frac{\partial \xi^r}{\partial t^\mu} = 0, \quad i = \overline{1,n}, \tag{2.6}$$

where $F^i = h^{\alpha\beta} F^{(i)}_{(\alpha)\beta}$.

To find other multi-time geometrical invariants for the PDE system (2.1), we also introduce the *$h-$KCC-covariant derivative of a $d-$tensor of type $\xi^i(t^\gamma)$* on the 1-jet space $J^1(T,M)$, via

$$\frac{\overset{h}{\nabla}\xi^i}{\partial t^\alpha} = \frac{\partial \xi^i}{\partial t^\alpha} + N^{(i)}_{(\alpha)r}\xi^r = \frac{\partial \xi^i}{\partial t^\alpha} + \frac{1}{2}\frac{\partial F^i}{\partial x^r_\gamma} h_{\gamma\alpha}\xi^r + \frac{1}{2} H^\gamma h_{\gamma\alpha}\xi^i,$$

where $H^\gamma = h^{\mu\nu} H^\gamma_{\mu\nu}$.

Remark 2.7 *The $h-$KCC-covariant derivative components $\dfrac{\overset{h}{\nabla}\xi^i}{\partial t^\alpha}$ transform under a change of coordinates (1.1) as a $d-$tensor of type $\mathrm{T}^{(i)}_{(\alpha)}$.*

In this geometrical context, the $h-$trace variational equations (2.6) can be rewritten in the following distinguished tensorial form:

$$h^{\alpha\beta}\frac{\overset{h}{\nabla}}{\partial t^\beta}\left[\frac{\overset{h}{\nabla}\xi^i}{\partial t^\alpha}\right] = \overset{h}{P}{}^i_r \xi^r,$$

where

$$\begin{aligned}\overset{h}{P}{}^i_j = & -\frac{\partial F^i}{\partial x^j} + \frac{1}{2}\frac{\partial^2 F^i}{\partial t^\gamma \partial x^j_\gamma} + \frac{1}{2}\frac{\partial^2 F^i}{\partial x^r \partial x^j_\gamma} x^r_\gamma - \frac{1}{2}\frac{\partial^2 F^i}{\partial x^j_\mu \partial x^r_\gamma} F^{(r)}_{(\gamma)\mu} + \\ & + \frac{h_{\gamma\mu}}{4}\frac{\partial F^i}{\partial x^r_\gamma}\frac{\partial F^r}{\partial x^j_\mu} + \frac{h^{\gamma\eta}}{2}\frac{\partial h_{\mu\gamma}}{\partial t^\eta}\frac{\partial F^i}{\partial x^j_\mu} + \\ & + \left[\frac{1}{2}\frac{\partial H^\gamma}{\partial t^\gamma} + \frac{h^{\gamma\eta}}{2}\frac{\partial h_{\mu\gamma}}{\partial t^\eta} H^\mu - \frac{h_{\gamma\mu}}{4} H^\gamma H^\mu\right]\delta^i_j.\end{aligned}$$

Definition 2.8 *The $d-$tensor $\overset{h}{P}{}^i_j$ is called the **second multi-time** $h-$**KCC-invariant** on the 1-jet space $J^1(T, M)$ of the PDE system (2.1), or the **multi-time** $h-$**deviation curvature** $d-$**tensor**.*

Example 2.9 *If we consider the second-order PDE system of the affine maps between the Riemannian manifolds $(T, h_{\alpha\beta}(t))$ and $(M, \varphi_{ij}(x))$, system which is given by (2.3), then the second multi-time $h-$KCC-invariant has the form*

$$\overset{h}{P}{}^i_j = -h^{\alpha\beta}\mathfrak{R}^i_{pqj} x^p_\alpha x^q_\beta,$$

where

$$\mathfrak{R}^i_{pqj} = \frac{\partial \gamma^i_{pq}}{\partial x^j} - \frac{\partial \gamma^i_{pj}}{\partial x^q} + \gamma^r_{pq}\gamma^i_{rj} - \gamma^r_{pj}\gamma^i_{rq}$$

*are the components of the curvature of the spatial Riemannian metric $\varphi_{ij}(x)$. Consequently, the $h-$trace variational equations (2.6) become the following **multi-time** $h-$**Jacobi field equations:***

$$h^{\alpha\beta}\left\{\frac{\overset{h}{\nabla}}{\partial t^\beta}\left[\frac{\overset{h}{\nabla}\xi^i}{\partial t^\alpha}\right] + \mathfrak{R}^i_{pqr} x^p_\alpha x^q_\beta \xi^r\right\} = 0,$$

where

$$\frac{\overset{h}{\nabla}\xi^i}{\partial t^\alpha} = \frac{\partial \xi^i}{\partial t^\alpha} + \gamma^i_{pr} x^p_\alpha \xi^r.$$

Example 2.10 *For the particular first order PDE system (2.4) the multi-time $h-$deviation curvature $d-$tensor is given by*

$$\begin{aligned}\overset{h}{P}{}^i_j = & \frac{h^{\alpha\beta}}{2}\left[\frac{\partial^2 X^{(i)}_{(\alpha)}}{\partial t^\beta \partial x^j} + \frac{\partial^2 X^{(i)}_{(\alpha)}}{\partial x^j \partial x^r} x^r_\beta + \frac{1}{2}\frac{\partial X^{(i)}_{(\alpha)}}{\partial x^r}\frac{\partial X^{(r)}_{(\beta)}}{\partial x^j}\right] + \\ & + \left[\frac{1}{2}\frac{\partial H^\gamma}{\partial t^\gamma} + \frac{h^{\gamma\eta}}{2}\frac{\partial h_{\mu\gamma}}{\partial t^\eta} H^\mu - \frac{h_{\gamma\mu}}{4} H^\gamma H^\mu\right]\delta^i_j.\end{aligned}$$

Definition 2.11 *The distinguished tensors*

$$\overset{h}{R}{}^{i\alpha}_{jk} = \frac{1}{3}\left[\frac{\partial \overset{h}{P}{}^{i}_{j}}{\partial x^{k}_{\alpha}} - \frac{\partial \overset{h}{P}{}^{i}_{k}}{\partial x^{j}_{\alpha}}\right], \qquad \overset{h}{B}{}^{i\alpha(\beta)}_{jk(l)} = \frac{\partial \overset{h}{R}{}^{i\alpha}_{jk}}{\partial x^{l}_{\beta}}$$

and

$$D^{(i)\ \ (\gamma)(\varepsilon)(\mu)}_{(\alpha)\beta(j)(k)(l)} = \frac{\partial^3 F^{(i)}_{(\alpha)\beta}}{\partial x^{j}_{\gamma}\partial x^{k}_{\varepsilon}\partial x^{l}_{\mu}}$$

are called the ***third**, **fourth*** *and* ***fifth multi-time*** $h-$***KCC-invariant*** *on the 1-jet vector bundle* $J^1(T,M)$ *of the PDE system (2.1).*

Remark 2.12 *Taking into account the transformation rules (2.2) of the components* $F^{(i)}_{(\alpha)\beta}$*, we immediately deduce that the components* $D^{(i)\ \ (\gamma)(\varepsilon)(\mu)}_{(\alpha)\beta(j)(k)(l)}$ *behave like a* $d-$*tensor on the 1-jet space* $J^1(T,M)$*.*

Example 2.13 *For the second-order PDE system (2.3) of the affine maps between the Riemannian manifolds* $(T, h_{\alpha\beta}(t))$ *and* $(M, \varphi_{ij}(x))$*, the third, fourth and fifth multi-time* $h-$*KCC-invariants are given by*

$$\overset{h}{R}{}^{i\alpha}_{jk} = h^{\alpha\mu}\mathfrak{R}^{i}_{pjk}x^{p}_{\mu}, \qquad \overset{h}{B}{}^{i\alpha(\beta)}_{jk(l)} = h^{\alpha\beta}\mathfrak{R}^{i}_{ljk}, \qquad D^{(i)\ \ (\gamma)(\varepsilon)(\mu)}_{(\alpha)\beta(j)(k)(l)} = 0.$$

Example 2.14 *For the first order PDE system (2.4) the third, fourth and fifth multi-time* $h-$*KCC-invariants are zero.*

Theorem 2.15 (of characterization of the multi-time KCC-invariants) *Let* (T,h) *be a Riemannian manifold, where* $m = \dim T \geq 3$*. If the first and the fifth KCC-invariants of the PDE system (2.1) cancel on* $J^1(T,M)$*, then there exist on* $J^1(T,M)$ *some local functions* $\Gamma^{i}_{pq}(t,x)$*, where* $i,p,q = \overline{1,n}$*,* $n = \dim M$*, and* $\mathtt{S}^{i\nu}_{\alpha pq}(t,x)$*,* $\alpha \neq \nu \in \{1,2,...,m\}$*,* $i = \overline{1,n}$*,* $p \neq q \in \{1,2,...,n\}$*, which have the properties*

$$\Gamma^{i}_{pq} = \Gamma^{i}_{qp}, \qquad \mathtt{S}^{i\nu}_{\alpha pq} + \mathtt{S}^{i\nu}_{\alpha qp} = 0$$

and (no sum by α *or* ν*)*

$$2\mathtt{S}^{i\nu}_{\alpha pq} = \sum_{\substack{\varepsilon=1\\ \varepsilon\neq\nu}}^{m}\left[h^{\varepsilon\varepsilon}\mathtt{S}^{i\nu}_{\varepsilon pq} - h^{\nu\nu}\mathtt{S}^{i\varepsilon}_{\nu pq}\right]h_{\varepsilon\alpha}, \tag{2.7}$$

such that (no sum by α *or* β*)*

$$F^{(i)}_{(\alpha)\beta} = \Gamma^{i}_{pq}x^{p}_{\alpha}x^{q}_{\beta} - H^{\mu}_{\alpha\beta}x^{i}_{\mu} + 2\delta_{\alpha\beta}\sum_{\substack{\nu=1\\ \nu\neq\alpha}}^{m}\ \sum_{p\neq q\in\{1,2,...,n\}}\mathtt{S}^{i\nu}_{\alpha pq}x^{p}_{\alpha}x^{q}_{\nu}, \tag{2.8}$$

where $\delta_{\alpha\beta}$ *is the Kronecker symbol and* $H^{\gamma}_{\alpha\beta}$ *are the Christoffel symbols of the Riemannian metric* $h_{\alpha\beta}(t)$*.*

Proof. By integration, the relations

$$D^{(i)\ \ (\gamma)(\varepsilon)(\sigma)}_{(\alpha)\beta(j)(k)(l)} = \frac{\partial^3 F^{(i)}_{(\alpha)\beta}}{\partial x^j_\gamma \partial x^k_\varepsilon \partial x^l_\sigma} = 0,$$

where $F^{(i)}_{(\alpha)\beta} = F^{(i)}_{(\beta)\alpha}$, subsequently lead to

$$\begin{aligned}
\frac{\partial^2 F^{(i)}_{(\alpha)\beta}}{\partial x^j_\gamma \partial x^k_\varepsilon} &= 2\Gamma^{(i)\ \ (\gamma)(\varepsilon)}_{(\alpha)\beta(j)(k)}(t,x) \Rightarrow \\
&\Rightarrow \ \frac{\partial F^{(i)}_{(\alpha)\beta}}{\partial x^j_\gamma} = 2\Gamma^{(i)\ \ (\gamma)(\nu)}_{(\alpha)\beta(j)(q)} x^q_\nu + \mathcal{U}^{(i)\ \ (\gamma)}_{(\alpha)\beta(j)}(t,x) \Rightarrow \\
&\Rightarrow \ F^{(i)}_{(\alpha)\beta} = \Gamma^{(i)\ \ (\mu)(\nu)}_{(\alpha)\beta(p)(q)} x^p_\mu x^q_\nu + \mathcal{U}^{(i)\ \ (\nu)}_{(\alpha)\beta(q)} x^q_\nu + \mathcal{V}^{(i)}_{(\alpha)\beta}(t,x),
\end{aligned}$$

where

$$\begin{aligned}
&\Gamma^{(i)\ \ (\gamma)(\varepsilon)}_{(\alpha)\beta(j)(k)} = \Gamma^{(i)\ \ (\gamma)(\varepsilon)}_{(\beta)\alpha(j)(k)}, \quad \Gamma^{(i)\ \ (\gamma)(\varepsilon)}_{(\alpha)\beta(j)(k)} = \Gamma^{(i)\ \ (\varepsilon)(\gamma)}_{(\alpha)\beta(k)(j)}, \\
&\mathcal{U}^{(i)\ \ (\gamma)}_{(\alpha)\beta(j)} = \mathcal{U}^{(i)\ \ (\gamma)}_{(\beta)\alpha(j)}, \qquad \mathcal{V}^{(i)}_{(\alpha)\beta} = \mathcal{V}^{(i)}_{(\beta)\alpha}.
\end{aligned} \tag{2.9}$$

The equalities $\overset{h}{\varepsilon}{}^{(i)}_{(\alpha)\beta} = 0$ on $J^1(T,M)$ lead us to

$$\begin{aligned}
\Gamma^{(i)\ \ (\mu)(\nu)}_{(\alpha)\beta(p)(q)} &= \frac{1}{2}\left[\Gamma^{i(\eta)(\nu)}_{(p)(q)}\delta^\mu_\beta + \Gamma^{i(\eta)(\mu)}_{(q)(p)}\delta^\nu_\beta\right] h_{\eta\alpha}, \\
\mathcal{U}^{(i)\ \ (\nu)}_{(\alpha)\beta(q)} &= \frac{1}{2}\mathcal{U}^{i(\eta)}_{(q)} h_{\eta\alpha}\delta^\nu_\beta + \frac{1}{2}H^\eta h_{\eta\alpha}\delta^\nu_\beta\delta^i_q - H^\nu_{\alpha\beta}\delta^i_q, \\
\mathcal{V}^{(i)}_{(\alpha)\beta} &= 0,
\end{aligned} \tag{2.10}$$

where

$$\Gamma^{i(\mu)(\nu)}_{(p)(q)} = h^{\varepsilon\rho}\Gamma^{(i)\ \ (\mu)(\nu)}_{(\varepsilon)\rho(p)(q)} \text{ and } \mathcal{U}^{i(\nu)}_{(q)} = h^{\varepsilon\rho}\mathcal{U}^{(i)\ \ (\nu)}_{(\varepsilon)\rho(q)}.$$

Applying an $h-$trace in the second relation of (2.10), we deduce that

$$\mathcal{U}^{(i)\ \ (\nu)}_{(\alpha)\beta(q)} = -H^\nu_{\alpha\beta}\delta^i_q.$$

The first relation of (2.10) and the first symmetry properties of (2.9) imply the following equalities:

1. for every $\alpha \neq \beta$ we have (no sum by α or β):
   (a) $\mu, \nu \notin \{\alpha, \beta\} \Rightarrow \Gamma^{(i)\ \ (\mu)(\nu)}_{(\alpha)\beta(p)(q)} = 0;$
   (b) $\mu = \alpha,\ \nu = \alpha \Rightarrow \Gamma^{(i)\ \ (\alpha)(\alpha)}_{(\alpha)\beta(p)(q)} = 0;$
   (c) $\mu = \alpha,\ \nu = \beta \Rightarrow \Gamma^{(i)\ \ (\alpha)(\beta)}_{(\alpha)\beta(p)(q)} = \dfrac{1}{2}\Gamma^{i(\eta)(\alpha)}_{(q)(p)} h_{\eta\alpha} \overset{not}{=} \mathbb{S}^i_{\alpha pq} = \mathbb{T}^i_{\beta pq};$
   (d) $\mu = \beta,\ \nu = \alpha \Rightarrow \Gamma^{(i)\ \ (\beta)(\alpha)}_{(\alpha)\beta(p)(q)} = \dfrac{1}{2}\Gamma^{i(\eta)(\alpha)}_{(p)(q)} h_{\eta\alpha} \overset{not}{=} \mathbb{T}^i_{\alpha pq} = \mathbb{S}^i_{\beta pq};$

(e) $\mu=\beta,\, \nu=\beta \Rightarrow \Gamma^{(i)\ \ (\beta)(\beta)}_{(\alpha)\beta(p)(q)} = \Gamma^{(i)\ \ (\beta)(\beta)}_{(\beta)\alpha(p)(q)} =$
$= \dfrac{1}{2}\left[\Gamma^{i(\eta)(\beta)}_{\ \ (p)(q)} h_{\eta\alpha} + \Gamma^{i(\eta)(\beta)}_{\ \ (q)(p)} h_{\eta\alpha}\right] = \mathtt{S}^{i\beta}_{\alpha pq} + \mathtt{T}^{i\beta}_{\alpha pq} = 0;$

2. for every $\alpha=\beta\in\{1,2,...,m\}$ we obtain (no sum by α):

   (a) $\mu\neq\alpha, \nu\neq\alpha \Rightarrow \Gamma^{(i)\ \ (\mu)(\nu)}_{(\alpha)\alpha(p)(q)} = 0;$

   (b) $\mu=\alpha, \nu\neq\alpha \Rightarrow \Gamma^{(i)\ \ (\alpha)(\nu)}_{(\alpha)\alpha(p)(q)} = \dfrac{1}{2}\Gamma^{i(\eta)(\nu)}_{\ \ (p)(q)} h_{\eta\alpha} \overset{not}{=} \mathtt{S}^{i\nu}_{\alpha pq} = \mathtt{T}^{i\nu}_{\alpha qp};$

   (c) $\mu\neq\alpha, \nu=\alpha \Rightarrow \Gamma^{(i)\ \ (\mu)(\alpha)}_{(\alpha)\alpha(p)(q)} = \dfrac{1}{2}\Gamma^{i(\eta)(\mu)}_{\ \ (q)(p)} h_{\eta\alpha} \overset{not}{=} \mathtt{T}^{i\mu}_{\alpha pq} = \mathtt{S}^{i\mu}_{\alpha qp};$

   (d) $\mu=\alpha,\, \nu=\alpha \Rightarrow \Gamma^{(i)\ \ (\alpha)(\alpha)}_{(\alpha)\alpha(p)(q)} = \Gamma^{(i)\ \ (\alpha)(\alpha)}_{(\alpha)\alpha(q)(p)} =$
   $= \dfrac{1}{2}\left[\Gamma^{i(\eta)(\alpha)}_{\ \ (p)(q)} h_{\eta\alpha} + \Gamma^{i(\eta)(\alpha)}_{\ \ (q)(p)} h_{\eta\alpha}\right] = \mathbb{T}^{i}_{\alpha pq} + \mathbb{S}^{i}_{\alpha pq};$

The first symmetry condition from (2.9), together with 1.(c) and 1.(d), give us ($m=\dim T\geq 3$)

$$\begin{array}{ccccccccccccc}
\mathbb{S}^i_{1pq} & = & \mathbb{T}^i_{2pq} & = & \mathbb{T}^i_{3pq} & = & \mathbb{T}^i_{4pq} & = & \cdot\ \ \cdot & = & \mathbb{T}^i_{mpq} & \overset{not}{=} & \dfrac{1}{2}\Gamma^i_{\ pq} \\
\mathbb{S}^i_{2pq} & = & \mathbb{T}^i_{1pq} & = & \mathbb{T}^i_{3pq} & = & \mathbb{T}^i_{4pq} & = & \cdot\ \ \cdot & = & \mathbb{T}^i_{mpq} & \overset{not}{=} & \dfrac{1}{2}\Gamma^i_{\ pq} \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot\ \ \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot\ \ \cdot & \cdot & \cdot & \cdot & \cdot \\
\mathbb{S}^i_{mpq} & = & \mathbb{T}^i_{1pq} & = & \mathbb{T}^i_{2pq} & = & \mathbb{T}^i_{3pq} & = & \cdot\ \ \cdot & = & \mathbb{T}^i_{(m-1)pq} & \overset{not}{=} & \dfrac{1}{2}\Gamma^i_{\ pq}.
\end{array}$$

Consequently, for every $\alpha\neq\beta\in\{1,2,...,m\}$ we have

$$\Gamma^{(i)\ \ (\mu)(\nu)}_{(\alpha)\beta(p)(q)} = \frac{1}{2}\Gamma^i_{\ pq}\left[\delta^\mu_\alpha\delta^\nu_\beta + \delta^\nu_\alpha\delta^\mu_\beta\right]$$

and for every $\alpha\in\{1,2,...,m\}$ we have

$$\Gamma^{(i)\ \ (\alpha)(\alpha)}_{(\alpha)\alpha(p)(q)} = \Gamma^i_{\ pq} = \Gamma^{(i)\ \ (\alpha)(\alpha)}_{(\alpha)\alpha(q)(p)} = \Gamma^i_{\ qp}.$$

Using now all the preceding properties, together with the equality 2.(b), we find the equations (2.7). Moreover, for every $\alpha\neq\nu\in\{1,2,...,m\}$, it is obvious that we have

$$\mathtt{S}^{i\nu}_{\alpha pq} + \mathtt{S}^{i\nu}_{\alpha qp} = 0 \Rightarrow \mathtt{S}^{i\nu}_{\alpha pp} = 0.$$

All the preceding situations can be briefly written in the general formula

$$\Gamma^{(i)\ \ (\mu)(\nu)}_{(\alpha)\beta(p)(q)} = \frac{1}{2}\Gamma^i_{\ pq}\left[\delta^\mu_\alpha\delta^\nu_\beta + \delta^\nu_\alpha\delta^\mu_\beta\right] + \mathtt{S}^{i\nu}_{\alpha pq}\delta_{\alpha\beta}\delta^\mu_\alpha\left[1-\delta^\nu_\alpha\right] + \mathtt{S}^{i\mu}_{\alpha qp}\delta_{\alpha\beta}\delta^\nu_\alpha\left[1-\delta^\mu_\alpha\right].$$

In conclusion, we obtain the equalities (2.8) on the 1-jet space $J^1(T,M)$. ■

Open problem. If we fix the indices i and $p \neq q$ in the set $\{1, 2, ..., n\}$, then we deduce that the system of equations (2.7) is an homogenous linear system of order $m(m-1)$. Consequently, it has at least the zero solution. Because the coefficients of the system depend only by the metric $h_{\alpha\beta}(t)$, there exist a temporal Riemannian metric $h_{\alpha\beta}(t)$ such that the system of equations (2.7) to admit only the banal solution?

References

[1] P.L. Antonelli, *Equivalence Problem for Systems of Second Order Ordinary Differential Equations*, Encyclopedia of Mathematics, Kluwer Academic, Dordrecht, 2000.

[2] V. Balan, M. Neagu, *Jet geometrical extension of the KCC-invariants*, http://arXiv.org/math.DG/0906.2903v2, (2009).

[3] V. Balan, I.R. Nicola, *Linear and structural stability of a cell division process model*, Hindawi Publishing Corporation, International Journal of Mathematics and Mathematical Sciences, vol. 2006, 1-15.

[4] E. Cartan, *Observations sur le mémoir précédent*, Mathematische Zeitschrift **37** (1933), no. **1**, 619-622.

[5] S.S. Chern, *Sur la géométrie d'un système d'equations differentialles du second ordre*, Bulletin des Sciences Mathématiques **63** (1939), 206-212.

[6] J. Eells, L. Lemaire, *A report on harmonic maps*, Bulletin of London Mathematical Society **10** (1978), 1-68.

[7] D.D. Kosambi, *Parallelism and path-spaces*, Mathematische Zeitschrift **37** (1933), no. **1**, 608-618.

[8] M. Neagu, *Riemann-Lagrange Geometry on 1-Jet Spaces*, Matrix Rom, Bucharest, 2005.

[9] M. Neagu, C. Udrişte, A. Oană, *Multi-time dependent sprays and h-traceless maps on $J^1(T, M)$*, Balkan Journal of Geometry and Its Applications, vol. **10** (2005), no. **2**, 76-92.

[10] V.S. Sabău, *Systems biology and deviation curvature tensor*, Nonlinear Analysis. Real World Applications **6** (2005), no. **3**, 563-587.

Authors' address:

Mircea NEAGU
University Transilvania of Braşov, Faculty of Mathematics and Informatics,
Department of Algebra, Geometry and Differential Equations,
B-dul Iuliu Maniu, Nr. 50, BV 500091, Braşov, Romania.
E-mail: mircea.neagu@unitbv.ro
Website: http://www.2collab.com/user:mirceaneagu